

Is guided self-help as effective as face-to-face psychotherapy for depression and anxiety disorders? A systematic review and meta-analysis of comparative outcome studies

P. Cuijpers^{1,2*}, T. Donker^{1,2}, A. van Straten^{1,2}, J. Li³ and G. Andersson^{4,5}

¹ Department of Clinical Psychology, VU University Amsterdam, The Netherlands

² EMGO Institute for Health and Care Research, VU University Amsterdam and VU University Medical Center, The Netherlands

³ Institute of Psychology, Chinese Academy of Sciences, Beijing, People's Republic of China

⁴ Department of Behavioural Sciences and Learning, Swedish Institute for Disability Research, Linköping University, Sweden

⁵ Department of Clinical Neuroscience, Psychiatry Section, Karolinska Institutet, Stockholm, Sweden

Background. Although guided self-help for depression and anxiety disorders has been examined in many studies, it is not clear whether it is equally effective as face-to-face treatments.

Method. We conducted a meta-analysis of randomized controlled trials in which the effects of guided self-help on depression and anxiety were compared directly with face-to-face psychotherapies for depression and anxiety disorders. A systematic search in bibliographical databases (PubMed, PsycINFO, EMBASE, Cochrane) resulted in 21 studies with 810 participants.

Results. The overall effect size indicating the difference between guided self-help and face-to-face psychotherapy at post-test was $d = -0.02$, in favour of guided self-help. At follow-up (up to 1 year) no significant difference was found either. No significant difference was found between the drop-out rates in the two treatments formats.

Conclusions. It seems safe to conclude that guided self-help and face-to-face treatments can have comparable effects. It is time to start thinking about implementation in routine care.

Received 12 December 2009; Revised 11 March 2010; Accepted 15 March 2010; First published online 21 April 2010

Key words: Bibliotherapy, depression, guided self-help, meta-analysis.

Introduction

The question whether depressed and anxious patients are capable of applying a cognitive behavioural intervention to themselves, with only minimal support from a professional therapist, has been examined for more than 40 years (Kahn & Baker, 1968; Hogan & Kirchner, 1968; Donner & Guerny, 1969; Watkins & Clum, 2008). Since then, dozens of randomized trials and meta-analyses have shown that guided self-help is effective in reducing depression, panic disorder, phobias and other anxiety disorders (Cuijpers, 1997; Gregory *et al.* 2004; Hirai & Clum, 2006; Gellatly *et al.* 2007; Menchola *et al.* 2007; Spek *et al.* 2007). Guided self-help can be defined as a psychological treatment, where the patient or client takes home a standardized

psychological treatment and works through it more or less independently (Marrs, 1995; Cuijpers & Schuurmans, 2007). In the standardized psychological treatment, the patient can use step-by-step instructions on how to apply a generally accepted psychological treatment procedure to himself. The standardized treatment can be written down in book form, or be made available through other media, such as the Internet, a stand-alone personal computer, television, video or audio. Guided self-help can be distinguished from other self-help interventions by the support that is given by a professional therapist or coach to the patient when working through the standardized treatment. The support given by the therapist should primarily be of supportive or facilitative nature, and is meant to support the patient in working through the standardized psychological treatment. Interaction between patient and therapist can take place through face-to-face contact, by telephone, by email, or any other communication method. An important distinction between guided self-help and

* Address for correspondence: P. Cuijpers, Ph.D., Professor of Clinical Psychology, Department of Clinical Psychology, VU University Amsterdam, Van der Boechorststraat 1, 1081 BT Amsterdam, The Netherlands.
(Email: p.cuijpers@psy.vu.nl)

face-to-face treatment is the amount of contact which is minimized in guided self-help.

Although a considerable number of studies have examined whether guided self-help is effective in the treatment of depression and anxiety disorders, it is not clear whether they are equally effective as face-to-face treatments. Most studies on guided self-help have found positive effects of these interventions compared with control conditions, and meta-analyses in this field have found large effect sizes (Gregory *et al.* 2004; Hirai & Clum, 2006; Gellatly *et al.* 2007; Menchola *et al.* 2007) which are comparable with those found for face-to-face treatments of depression and anxiety. However, these meta-analyses may very well be influenced by factors that differed among the various studies, such as length of treatment, type of treatment, or initial symptom severity (Shadish & Sweeney, 1991; Spielmans *et al.* 2007). Hence, possible differences between the effects of the two treatment formats may very well be artifacts, which do not reflect true superiority of one of the types of treatment over the other (Spielmans *et al.* 2007). Direct comparisons of guided self-help and face-to-face treatments, in which patients are assigned to one of the two treatments in the same study, are better equipped to rule out the influence of study characteristics, and they certainly provide more reliable evidence about a possible superiority of one type of therapy over the other (Spielmans *et al.* 2007).

Although a considerable number of studies have compared guided self-help and face-to-face treatment directly with each other (i.e. in the same study), no meta-analysis of these studies has been conducted. Such a meta-analysis is, however, important, because most individual studies in this field do not have sufficient statistical power to detect a significant difference between the two treatments. In fact, many studies could be better described as equivalence studies in which a null finding is the expected outcome (Piaggio *et al.* 2006). If there is a difference between the two treatments it can be assumed that this difference is relatively small, and in order to detect a small effect, a large number of participants is needed. Most studies in this field have not included sufficient participants to detect a possible difference. This could suggest that the two treatments are equally effective, while in fact they are not. A meta-analysis can solve this problem because the results of multiple studies are combined, and sufficient statistical power is available to detect a possible difference.

One recent meta-analysis of studies examining self-help interventions for anxiety disorders has included a subset of studies in which guided self-help was compared directly with face-to-face treatments for anxiety disorders (Hirai & Clum, 2006). This meta-analysis

did not find a significant advantage of face-to-face treatment over guided self-help (effect size $d = 0.11$), or the other way around. This meta-analysis included, however, only a selection of the currently available studies (nine of the 15 studies on anxiety disorders were included in the current meta-analysis). Furthermore, no power calculation was conducted to assess whether the available studies had sufficient power to detect a significant effect. Finally, sources of heterogeneity were not examined, and nor was publication bias.

We decided to conduct a new meta-analysis of randomized controlled trials in which the effects of guided self-help on depression and anxiety were compared directly with face-to-face psychotherapies for depression and anxiety disorders.

Method

Identification and selection of studies

We used several methods to identify studies for inclusion. First, we used a database of 1036 papers on the psychological treatment of depression, which includes studies comparing guided self-help and face-to-face therapies. This database has been described in detail elsewhere (Cuijpers *et al.* 2008) and has been used in a series of earlier meta-analyses (www.evidencebasedpsychotherapies.org). It was developed through a comprehensive literature search (from 1966 to January 2009) in which we examined 9011 abstracts in PubMed (1629 abstracts), PsycINFO (2439), EMBASE (2606) and the Cochrane Central Register of Controlled Trials (2337). These abstracts were identified by combining terms indicative of psychological treatment and depression [both medical subject heading (MeSH) terms and text words]. For this database, we also collected the primary studies from 42 meta-analyses of psychological treatment for depression (www.evidencebasedpsychotherapies.org). Second, we conducted additional searches in bibliographical databases, in which we combined search terms indicative of guided self-help and each of the disorders we examined, and randomized controlled trials as a limit. An example of a search string can be found in Supplementary Appendix B of this paper (available online).

Third, we examined the reference lists of all meta-analyses examining guided self-help interventions for depression and anxiety mentioned earlier. Fourth, we checked the references of the included primary studies. We did not contact study authors for additional data, unpublished studies and studies in press.

We included randomized studies in which guided self-help was compared with a face-to-face treatment

for depression or anxiety, and which reported outcomes on depression and/or anxiety. Guided self-help was defined as a treatment (1) in which the procedures were written down (or presented in an audio or video file) and (2) the patients had to work through these procedures more or less independently, while (3) the therapist gave support in working through the procedures (4) with limited contact. Limited contact could be delivered during brief personal contacts, by telephone or by email. We allowed for a maximum of 12 contacts, with a maximum of 20 min each. Face-to-face treatments had to use the same format and contents as the guided self-help, but also had to use full individual or group treatment sessions to deliver the treatment.

We included studies in which a diagnostic interview was used to establish the presence of depression or anxiety disorders, but we also included studies which used other inclusion criteria. We excluded studies in children and adolescents, as well as studies examining virtual reality treatments (Côté & Bouchard, 2008). No language restrictions were applied.

A detailed list of the variables we extracted from each study can be found in Supplementary Appendix C (available online). Data abstraction from the studies was conducted by the first author (P.C.) and checked by the third author (A.v.S.).

Quality assessment

We assessed the validity of included studies using four criteria of the 'risk of bias' assessment tool, developed by the Cochrane Collaboration (Higgins & Green, 2008). This tool assesses possible sources of bias in randomized trials, including the adequate generation of allocation sequence; the concealment of allocation to conditions; the prevention of knowledge of the allocated intervention; and dealing with incomplete outcome data. The two other criteria of the 'risk of bias' assessment tool were not used in this study, because we found no clear indication in any of the studies that these had influenced the validity of the study (suggestions of selective outcome reporting; and other problems that could put it at a high risk of bias).

We also rated the quality of the treatment implementation using three criteria which were based on an authoritative review of empirically supported psychotherapies (Chambless & Hollon, 1998): (1) the study referred to the use of a treatment manual (either a published manual, or a manual specifically designed for the study); (2) the therapists who conducted the therapy were trained for the specific therapy, either specifically for the study or as a general training; (3) treatment integrity was checked during the study

(by supervision of the therapists during treatment or by recording of treatment sessions or by systematic screening of protocol adherence by a standardized measurement instrument).

Meta-analyses

For each comparison between guided self-help and a face-to-face treatment, we calculated the effect size indicating the difference between the two groups at post-test (Cohen's *d*). Effect sizes were calculated by subtracting (at post-test) the average score of the guided self-help group from the average score of the face-to-face treatment group, and dividing the result by the pooled standard deviations of the two groups. Effect sizes of 0.56 to 1.2 can be assumed to be large, effect sizes of 0.33 to 0.55 are moderate, and effect sizes of 0 to 0.32 are small (Lipsey, 1990).

In the calculations of effect sizes we only used those instruments that explicitly measured symptoms of depression, and in the studies that examined anxiety, instruments that explicitly measured symptoms of anxiety. If more than one measure was used, the mean of the effect sizes was calculated, so that each study provided only one effect size. If means and standard deviations were not reported, we used the procedures of the COMPREHENSIVE META-ANALYSIS software (see below; Biostat, Inc., USA) to calculate the effect size using dichotomous outcomes. In two studies, it was only reported that there was no significant difference between the two conditions (Baker *et al.* 1973; Ghosh *et al.* 1988). In these studies, the effect size was assumed to be zero.

To calculate pooled mean effect sizes, we used the computer program COMPREHENSIVE META-ANALYSIS (version 2.2.021; Biostat, Inc., USA). As we expected considerable heterogeneity among the studies, we decided to calculate mean effect sizes using a random-effects model.

As a test of homogeneity of effect sizes, we calculated the I^2 statistic, which is an indicator of heterogeneity in percentages. A value of 0% indicates no observed heterogeneity, and larger values show increasing heterogeneity, with 25% as low, 50% as moderate, and 75% as high heterogeneity (Higgins *et al.* 2003). We also calculated the Q statistic, but only report whether this was significant or not.

Publication bias was tested by inspecting the funnel plot on primary outcome measures, and by Duval & Tweedie's trim and fill procedure (Duval & Tweedie, 2000) which yields an estimate of the effect size after the publication bias has been taken into account (as implemented in COMPREHENSIVE META-ANALYSIS, version 2.2.021).

Apart from the outcomes on depression and anxiety, we also calculated the relative risk of dropping out from the treatments. Again, we conducted all meta-analyses with the random-effects model and we calculated the Q statistic and the I^2 statistic to estimate heterogeneity between study outcomes.

In order to examine possible predictors of the difference between guided self-help and face-to-face treatment, we conducted a multivariate meta-regression analysis with the effect size as the dependent variable. As predictors, we used the characteristics of the populations (disorder: depression, panic disorder, phobia; community *versus* other type of recruitment; whether or not a diagnostic interview was conducted; aimed at adults in general or a more specific target group); the interventions (personal contact support *versus* other support during guided self-help; book *versus* other medium; individual *versus* group treatment) and the quality of the studies (met more than three quality criteria). In order to avoid collinearity among the predictors that were entered in the regression model, we first examined whether high correlations were found among the variables that could be entered into the model. The correlations between all entered characteristics were calculated. We found that none of the correlations was higher than 0.50. The multivariate meta-regression analyses were conducted in Stata SE/8 for Windows (StataCorp LP, USA).

Power calculation

We assumed the two treatment formats to be equally effective if the differential effect size was small. Although there is no clear agreement on what should be considered to be a small effect size, we used the definition of Lipsey (1990), which says that effect sizes of 0.3 can be considered as small. To have sufficient statistical power in our meta-analysis to be able to detect a small effect size, we conducted a power calculation according to the procedures described by Borenstein *et al.* (2009). These calculations indicated that we would need to include at least 14 studies with a mean sample size of 50 (25 participants per condition), to be able to detect an effect size of $d=0.3$ (conservatively assuming a high level of between-study variance, τ^2 , a statistical power of 0.80, and a significance level, α , of 0.05). Alternatively, we would need 18 studies with 40 participants each to detect an effect size of $d=0.30$, or 24 studies with 30 participants.

Results

Selection and inclusion of studies

We examined a total of 2761 abstracts, and 91 publications were retrieved for possible inclusion. We




Fig. 1. Flowchart of inclusion of studies. GSH, Guided self-help; FTF, face-to-face psychotherapy.

excluded 70 studies, because no guided self-help was examined (33 studies), because guided self-help was not compared with a face-to-face treatment (17 studies), because the studies examined virtual reality (nine studies), because the studies were not randomized trials (four studies), because they examined children or adolescents (three studies), or because of other reasons (four studies). The remaining 21 studies met all inclusion criteria and were included in our meta-analysis. A flowchart describing the inclusion of studies is presented in Fig. 1.

Characteristics of included studies

The 21 studies included a total of 810 participants (429 in face-to-face conditions and 381 in the guided self-help conditions). Selected characteristics of the included studies are presented in Table 1.

Most studies (14 out of 21) were aimed at adults in general, five were (largely) aimed at student populations, and two were aimed at more specific populations (one at older adults, and one at adults with co-morbid substance-use problems). Of the studies, six studies were aimed at patients with depression, seven at panic, three at social phobia, two at specific phobias, and three at phobias in general. In 15 of the 21 studies, a diagnostic interview was used to establish the presence of the depressive or anxiety disorder. Patients were recruited from the general population through media announcements in 17 studies, while three studies recruited people from clinical populations (one study did not report the recruitment method).

In the 21 studies, a total of 24 comparisons were made between a guided self-help condition and a face-to-face therapy. All treatments were cognitive and

Table 1. Selected characteristics of studies comparing face-to-face therapies with guided self-help for depression or anxiety disorders

Study	Disorder	Target group	Diagnosis	Recruitment	FTF therapy	Format	No. of sessions	n	GSH	Type	n	Instruments	Country	Quality ^a						
														a	b	c	d	e	f	g
Baker <i>et al.</i> (1973)	Acrophobia	Adults	–	Comm	Relaxation, exposure	Ind	13	8	One FTF session + tape-recorded sessions + six therapist meetings (5 min)	Audio	9	AQ	USA	–	–	–	–	–	+	–
Brown & Lewinsohn (1984)	Depressive disorder	Adults	RDC	Comm	Cognitive restructuring, activity scheduling, social skills – individual As above – group	Ind	12	13	One FTF session + 11 telephone contacts (20 min)	Book	14	BDI, CES-D	USA	–	–	±	–	+	+	+
Carlbring <i>et al.</i> (2005)	Panic disorder w/wo agoraphobia	Adults	CIDI, ADIS, SCID	Comm	Breathing retraining, cognitive restructuring, exposure, assertiveness training	Grp	12	25	Internet + emails	Internet	14 25	BSQ, MI, BAI	SW	+	+	±	+	+	+	+
Floyd <i>et al.</i> (2004)	Depression (MDD, DYS, minD)	Older adults	DSM-IV	Comm	CBT (according to Beck <i>et al.</i> 1979)	Ind	12–20	16	Book + four telephone calls	Book	16	HAMD, GDS	USA	–	–	+	+	+	+	+
Ghosh <i>et al.</i> (1988)	Phobias	Adults	ICD-9	Clin	Exposure	Ind	7	19	Computer-instructed exposure + 5 min therapist contact	Comp	28	FQ	UK	–	–	+	–	–	+	–
Gould <i>et al.</i> (1993)	Panic disorder w/wo agoraphobia	Adults (mostly students)	ADIS-R	Comm	Exposure, relaxation, cognitive restructuring, breathing retraining	Ind	8	9	Book + two telephone calls (10 min)	Book	12	DPAR, PACQ, MI	USA	–	–	±	–	+	–	+
Hecker <i>et al.</i> (1996)	Panic disorder w/wo agoraphobia	Adults	ADIS-R, SCID-II	NR	Breathing retraining, cognitive restructuring, exposure	Ind	12	7	Book + four therapist contacts	Book	5	Number of panic attacks	USA	–	–	–	–	+	–	+
Hecker <i>et al.</i> (2004)	Panic disorder w/wo agoraphobia	Adults	ADIS-IV	Comm	Breathing retraining, cognitive restructuring, exposure	Grp	4	31	One FTF session + three telephone contacts	Book	17	Clinical sign change	USA	–	–	+	–	+	–	–
Kahn & Baker (1968)	Phobias	Students	–	Comm	Exposure	Ind	12	7	One FTF session + six telephone contacts	Audio	6	Improvement rate	USA	–	–	–	–	+	–	+

[continued overleaf]

Selmi <i>et al.</i> (1990)	Depression (MDD, minD)	Adults	RDC	Comm	Cognitive restructuring, activity scheduling	Ind	6	12	Computer program + brief help at start up and end	Comp	12	BDI, SCL-90-d, HAMD	USA	-	-	+	+	+	+	-
Sharp <i>et al.</i> (2000)	Panic disorder w/wo agoraphobia	Adults	Diagnosis according to DSM-III-R	Clin	Cognitive behavioural intervention	Ind	8	31	Book + six brief sessions	Book	31	GSS	UK	+	+	+	-	+	+	-
Tillfors <i>et al.</i> (2008)	Social phobia and public speaking fear	University students	SPSQ	Comm	Cognitive restructuring, exposure, social skills, relapse prevention	Grp	5	19	Internet + emails	Internet	19	BAI, LSAS, SIAS, SPS, SPSQ	SW	-	-	±	+	+	+	+
Vestre & Judge (1989)	Social anxiety	University students	-	Comm	Rational emotive therapy	Grp	5	20	Book + five telephone calls	Book	21	FNE, HSCL-a, SAD	USA	-	-	±	-	+	+	-
Wollersheim & Wilson (1991)	Depression	Adults	Diagnosis according to DSM-III	Comm	Cognitive restructuring, activity scheduling, relaxation, problem-solving	Grp	10	8	Book + three sessions	Book	8	BDI, CR, MMPI-d, SR	USA	-	-	+	-	+	-	+

ADIS, Anxiety Disorders Interview Schedule; ADIS-IV, Anxiety Disorders Interview Schedule for DSM-IV; ADIS-R, Anxiety Disorders Interview Schedule Revised; AQ, Acrophobia Questionnaire; AU, Australia; BA-f, frequency of panic attacks; BA-g, goals (blind assessor); BA-mp, main problem (blind assessor); BAI, Beck Anxiety Inventory; BAT, Behavior Approach Test; BCL; Timed Behavioral Checklist for Performance Anxiety; BDI, Beck Depression Inventory; BDI-II, Beck Depression Inventory, 1996 Revision; BSQ, Body Sensations Questionnaire; CA, Canada; CBT, cognitive behaviour therapy; CES-D, Center for Epidemiological Studies – Depression scale; CIDI, Composite International Diagnostic Interview; Clin, recruitment from clinical samples; Comm, community recruitment; Comp, computer; CR, clinician rating of depression; CR-a, clinician-rated agoraphobia; CR-p, clinician-rated panic; DASS, Depression, Anxiety, Stress Scales; DPAR, Daily Panic Attack Record; DSM-III, Diagnostic and Statistical Manual of Mental Disorders, third edition; DSM-III-R, Diagnostic and Statistical Manual of Mental Disorders, third edition revised; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, fourth edition; DYS, dysthymia; FNE, Fear of Negative Evaluation Scale; FQ, Fear Questionnaire; FSS, Fear Survey Schedule; FT, Fear Thermometer; FTF, face-to-face therapy; GDS, Geriatric Depression Scale; Grp, group; GSH, guided self-help; GSS, global symptom severity; HAMD, Hamilton Rating Scale of Depression; HSCL-a, Hopkins Symptom Checklist, anxiety subscale; ICD-9, International Classification of Diseases, 9th revision; Ind, individual; LSAS, Liebowitz Social Anxiety Scale self-report version; MDD, major depressive disorder; MI, Mobility Inventory; minD, minor depression; MMPI, Minnesota Multiphasic Personality Inventory; MMPI-d; Minnesota Multiphasic Personality Inventory Depression Scale; NR, not reported; PACQ, Panic Attack Cognitions Questionnaire; PASQ, Panic Attack Symptoms Questionnaire; PDSS, Panic Disorder Severity Scale; POMS, Profile of Mood States; RDC, Research Diagnostic Criteria; SAD, Social Avoidance and Distress Scale; SCID, Structured Clinical Interview for DSM Disorders; SCID-II, Structured Clinical Interview for DSM-IV Axis II Personality Disorders; SCID-RV, Structured Clinical Interview for DSM Disorders Research Version; SCL-90-d, Symptom Checklist-90 depression scale; SDS, Self-rating Depression Scale; SIAS, Social Interaction Anxiety Scale; SNAQ, Snake Attitude Questionnaire; SPS, Social Phobia Scale; SPSQ, Social Phobia Screening Questionnaire; SR, self-rating of depression; SR-g, goals (self-rated); SR-mp, main problem (self-rated); SUD, Subjective Units of Disturbance; SW, Sweden; w/wo, with or without.

^a Quality assessment: a, allocation sequence adequately generated; b, allocation adequately concealed; c, knowledge of the allocated interventions adequately prevented (blinding); ± indicates that only self-report measures were used; d, incomplete outcome data adequately addressed; e, manual available; f, therapist trained; g, integrity check of intervention.

Table 2. Meta-analyses of studies comparing the effects of guided self-help and face-to-face psychotherapies for adult depression and anxiety disorders

	No. of studies	<i>d</i>	95% CI	Z	<i>I</i> ^{2a}
Effect sizes at post-test					
All comparisons	24	-0.02	-0.20 to 0.15	-0.24 N.S.	26.34 N.S.
Outlier excluded ^b	23	0.02	-0.14 to 0.18	0.29 N.S.	12.19 N.S.
One ES per study (highest) ^c	21	0.03	-0.15 to 0.21	0.34 N.S.	22.88 N.S.
One ES per study (lowest) ^c	21	-0.03	-0.23 to 0.17	-0.29 N.S.	34.93 N.S.
Effect sizes at follow-up					
1-3 months	10	-0.06	-0.30 to 0.17	-0.52 N.S.	0 N.S.
4-6 months	9	0.08	-0.17 to 0.33	0.61 N.S.	0 N.S.
12 months	3	-0.27	-0.62 to 0.07	-1.55 N.S.	0 N.S.

CI, confidence interval; N.S., non-significant; ES, effect size.

^a The *p* values in this column indicate whether the *Q* statistic is significant (the *I*² statistics does not include a test of significance).

^b Hecker *et al.* (2004).

^c In these analyses only one comparison from each study was used.

behavioural in nature. In the 24 comparisons, 16 of the face-to-face treatments used an individual treatment format while the remaining eight used a group treatment format. The number of treatment sessions in the face-to-face treatments ranged from four to 16. In the 24 guided self-help conditions, 15 used a self-help book, four a stand-alone computer program, three used an Internet-based intervention, and two used audio recordings to deliver the treatment. Support was given by brief personal contact in five comparisons, brief personal contact plus telephone calls in seven comparisons, by telephone calls only in five comparisons, and by email in three comparisons. In the remaining four comparisons the patient worked alone on a personal computer at the therapist's office and received brief face-to-face support.

Of the studies, 13 studies were conducted in the USA, three in the UK, and the remaining five in other countries (Australia, Sweden, Canada).

Quality of included studies

The quality of the studies was not optimal. Of the 21 studies, 16 gave insufficient information whether the allocation sequence was generated adequately. Also, 17 studies gave insufficient information about whether the allocation was adequately concealed. We assessed whether incomplete outcome data were adequately addressed, by conducting intention-to-treat analyses with all randomized subjects being included in the analyses. This was the case in only five studies. However, in 18 studies knowledge of the allocated interventions was adequately prevented by blinding of the assessors or because only self-report measures

were used (and blinding of assessors was not relevant). Only one study met all of the four quality criteria (Carlbring *et al.* 2005), and four studies met three of the four criteria (Sharp *et al.* 2000; Marks *et al.* 2004; Kiroopoulos *et al.* 2008; Kay-Lambkin *et al.* 2009).

The quality of the treatment implementation was good in most studies. In 19 studies a treatment manual was used, in 15 studies the therapists were specifically trained for the intervention, and in 12 studies an integrity check of the interventions was conducted. In seven studies all three criteria for the quality of treatment implementation were met (Rosen *et al.* 1976; Schmidt & Miller, 1983; Brown & Lewinsohn, 1984; Floyd *et al.* 2004; Carlbring *et al.* 2005; Tillfors *et al.* 2008; Kay-Lambkin *et al.* 2009).

Differences between guided self-help and face-to-face treatments

The mean effect size indicating the difference between guided self-help and face-to-face psychotherapy was $d = -0.02$ [95% confidence interval (CI) -0.20 to 0.15, N.S.], in favour of guided self-help. Heterogeneity was low ($I^2 = 26.34$) and not statistically significant (Table 2). The effect sizes and 95% CIs of the studies are plotted in Fig. 2. Our sample of studies had sufficient statistical power to detect a differential effect size of $d = 0.30$.

The 95% CI of the effect size found for one study (Hecker *et al.* 2004) did not overlap with the CI of the pooled effect size, and may be an outlier. However, removal of this study hardly affected the outcome ($d = 0.02$, 95% CI 0.14-0.18, N.S., $I^2 = 12.19$, N.S.).




Fig. 2. Differences between guided self-help (GSH) and face-to-face (FTF) treatments for depression and anxiety disorders: standardized effect sizes. Std diff, Standardized difference; CI, confidence interval; I, individual; G, group; sG, small group; IG, large group.

We also examined the influence of the two studies in which the effect size was assumed to be zero because no significant difference was found between guided self-help and face-to-face therapy, by removing them from the analyses. The results of these analyses, however, were virtually the same as the overall analyses ($d = -0.01$, 95% CI -0.20 to 0.19 , $I^2 = 26.34$).

In this meta-analysis we included two studies in which more than one comparison was made between guided self-help and a face-to-face treatment (Schmidt & Miller, 1983; Brown & Lewinsohn, 1984). This means that multiple comparisons from these two studies were included in the same analysis. These multiple comparisons, however, are not independent of each other, which may have resulted in an artificial reduction of heterogeneity and may have affected the pooled effect size. We examined the possible effects of this in two different ways. First, we divided the number of respondent over the different comparison groups in the studies with multiple comparison groups, so that we included each respondent only once in the overall analyses. The results of these analyses are almost the same as in our main analyses ($d = 0.00$, 95% CI -0.17 to 0.18 , $I^2 = 19.35$). In our second approach, we included only one comparison per study in the analysis (Table 2). First we conducted an analysis in which we

included only one effect size per study and chose the one with the largest effect size from the studies with multiple comparisons. Then we conducted another analysis in which we included only the smallest effect size. As can be seen from Table 2, the resulting effect sizes were almost the same as in the overall analyses. Heterogeneity did not increase very much, and was not statistically significant in any analysis.

In our analyses, we included studies with more than one outcome measure and we calculated the mean of the effect sizes based on these outcome measures. However, this approach assumes that the correlation between the measures within the same study is 1.00, which is not necessarily the case. Therefore we conducted another meta-analysis in which we assumed that all effect sizes within one study were independent (and the correlation would be zero). In Appendix A we have presented the forest plot of these analyses. The resulting overall effect size was almost the same as found in the main analyses ($d = -0.03$ in favour of guided self-help, with very low heterogeneity, $I^2 = 12.85$).

In order to examine the influence of individual studies, we examined which study had the largest impact on the overall effect size. Removal of the study by Sharp *et al.* (2000) resulted in the largest decrease of the effect size (the resulting effect size was

Table 3. Regression coefficients of study characteristics in relation to the effect size of guided self-help versus face-to-face treatments of depression: multivariate meta-regression analyses

	B	95% CI	p
Disorder			
Depression	Reference		
Panic	0.19	−0.63 to 1.01	0.65
Phobias	0.27	−0.69 to 1.22	0.58
Aimed at adults in general (yes/no)	−0.18	−0.75 to 0.39	0.53
Community recruitment (yes/no)	−0.26	−1.14 to 0.63	0.57
Diagnostic interview (yes/no)	−0.10	−0.77 to 0.57	0.78
Support: personal contact (yes/no)	0.01	−0.71 to 0.74	0.97
Book or other medium	0.21	−0.37 to 0.79	0.49
Number of sessions (continuous)	−0.01	−0.10 to 0.07	0.73
Individual format (yes/no)	0.38	−0.22 to 0.98	0.21
Met > three quality criteria (yes/no)	0.32	−0.23 to 0.87	0.26
Constant	−0.24	−2.58 to 2.10	0.84

CI, Confidence interval.

$d = -0.06$). After the removal of this study, we repeated this procedure and examined which study should be removed in order to realize the next largest decrease of the effect size. This was the study by Kiroopoulos *et al.* (2008), and the meta-analysis resulted in an effect size of $d = -0.09$. Repeating this procedure a third time resulted (after removal of the study by Kay-Lambkin *et al.* 2009) in a mean effect size of $d = -0.12$. None of the three resulting effect sizes were significantly different from zero. A comparable procedure to examine whether individual studies resulted in an increase of the effect size (in favour of face-to-face treatment) indicated that removal of the study by Hecker *et al.* (2004) resulted in the largest increase (resulting effect size $d = 0.02$), followed by the study of Schmidt *et al.* (1983) (effect size $d = 0.06$), and the study by Floyd *et al.* (2004) (effect size $d = 0.08$). These analyses did not suggest that removal of individual studies resulted in major changes in the overall effect size.

We could calculate effect sizes indicating the difference between face-to-face therapies at follow-up in 17 studies. We divided the effect sizes according to the follow-up period (1–3 months, 4–6 months, 12 months). As can be seen in Table 2, none of the effect sizes differed significantly from zero.

Neither the funnel plot nor Duval & Tweedie's trim and fill procedure pointed to a significant publication bias. The effect size indicating the difference in reduction of depressive symptomatology between guided self-help and face-to-face treatments did not change after adjustment for possible publication bias (the observed and adjusted effect sizes were exactly

the same, and the number of imputed studies was zero).

We examined possible predictors of the difference between guided self-help and face-to-face treatment, with multivariate meta-regression analysis. The results of these analyses are presented in Table 3. As can be seen, none of the predictors was significantly associated with the effect size.

Drop-out

The definition of drop-out differed considerably between studies, ranging from drop-out from the interventions to drop-out from the studies, and many variations in between. In 18 studies (21 comparisons) data on drop-out were presented. We calculated the relative risk of dropping out from guided self-help and face-to-face treatments in these studies. One study reported no drop-out in both conditions, and was not included in these analyses (Selmi *et al.* 1990). The pooled relative risk of dropping out was 1.14 (95% CI 0.77–1.67), indicating that the drop-out rate was somewhat higher in the guided self-help conditions, but that was not significant ($p = 0.52$). Heterogeneity (I^2) was zero and not significant ($Q = 17.05$, $p > 0.1$).

Discussion

In this study, we found no indication that the effects of guided self-help and face-to-face treatments differ significantly from each other, although we had sufficient statistical power to detect small differences. This was also true at follow-up periods of up to 1 year.

Furthermore, we found no indication that drop-out rates differed between the two treatment formats.

Although this meta-analysis found support for the hypothesis that guided self-help and face-to-face treatments for depression and anxiety do not significantly differ in effectiveness, this does not imply that guided self-help is effective in all patients seeking help in mental health care or primary care. The studies examined in this meta-analysis only included patients who were willing to be randomized to both conditions. People who are not interested in guided self-help have probably not participated in these trials. This means that guided self-help and face-to-face treatments may indeed be equally effective for many, but not for all people with mood or anxiety disorders. Future research should be conducted to examine who is willing to participate in guided self-help treatments and who is not, and if there are differential predictors and mediators of outcome (Andersson *et al.* 2008).

Most research included in this meta-analysis was conducted with people who are recruited from the community by media announcements. More research is needed to examine how guided self-help can be used in clinical practice. There are some indications that patients referred to guided self-help by their general practitioner benefit most from these treatments, compared with patients who are referred by mental health professionals and self-referrals from the community, while self-referred patients improve more than patients referred by mental health professionals (Mataix-Cols *et al.* 2006).

An interesting issue that results from this study is that apparently the patient–therapist relationship can be realized with minimal contact with the therapist. This may suggest that it is not so much the intensity of the contact that makes a well-functioning relationship possible (Knaevelsrud & Maercker, 2007), but more the contact between the two in itself. It is also possible that the patient–therapist relationship is not needed at all in these treatments, although there is evidence that unguided self-help is significantly less effective than guided self-help, both in depression and anxiety disorders (Hirai & Clum, 2006; Spek *et al.* 2007). This study has several limitations. One important limitation is the relatively small sample of studies, which limits the possibility to explore differences between subgroups. For example, we did not have sufficient power to detect a small differential effect size for the whole sample of studies and for the studies on anxiety disorders, but not for the studies on depression. Second, the quality of the included studies was not optimal in many studies. Third, we may have missed studies that were unpublished or in the press. Fourth, we included a broad range of disorders in our meta-analysis. Depression and anxiety disorders are

different categories of mental disorders and within the category of anxiety disorders, we included studies focusing on panic disorders, phobias in general and specific categories of phobias. However, we found very low levels of heterogeneity, suggesting that the different diagnostic categories did not result in different answers to the question whether the two treatments are equally effective. Furthermore, co-morbidity between anxiety disorders and depression is very high, suggesting that they are closely related to each other, and cognitive and behavioural treatment strategies for depression and anxiety share many common elements.

Despite these limitations, it seems safe to conclude that guided self-help and face-to-face treatments for depression and anxiety have comparable effects, and that there is no evidence that one or the other is significantly larger than the other. There is no reason not to consider using guided self-help as a complement in clinical practice, and we suspect that face-to-face treatment and guided self-help will blend in with each other increasingly in the near future, for example by using computer assistance when providing psychoeducational material (Craske *et al.* 2009).

Note

Supplementary material accompanies this paper on the Journal's website (<http://journals.cambridge.org/psm>).

Acknowledgements

None.

Declaration of Interest

None.


References

- Andersson G, Carlbring P, Grimlund A (2008). Predicting treatment outcome in Internet *versus* face to face treatment of panic disorder. *Computers in Human Behavior* **24**, 1790–1801.
- Baker BL, Cohen DC, Saunders JT (1973). Self-directed desensitization for acrophobia. *Behaviour Research and Therapy* **11**, 79–89.
- Beck AT, Rush AJ, Shaw BF, Emery G (1979). *Cognitive Therapy of Depression*. Guilford Press: New York.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009). *Introduction to Meta-Analysis*. Wiley: Chichester, UK.
- Brown RA, Lewinsohn PM (1984). A psychoeducational approach to the treatment of depression: comparison of group, individual and minimal contact procedures.

- Journal of Consulting and Clinical Psychology* **52**, 774–783.
- Carlbring P, Nilsson-Ihrfelt E, Waara J, Kollenstam C, Buhrman M, Kaldø V, Söderberg M, Ekselius L, Andersson G** (2005). Treatment of panic disorder: live therapy vs. self-help via Internet. *Behaviour Research and Therapy* **43**, 1321–1333.
- Chambless DL, Hollon SD** (1998). Defining empirically supported therapies. *Journal of Consulting and Clinical Psychology* **66**, 7–18.
- Côté S, Bouchard S** (2008). Virtual reality exposure for phobias: a critical review. *Journal of Cyber Therapy and Rehabilitation* **1**, 75–91.
- Craske MG, Rose RD, Lang A, Welch SS, Campbell-Sills L, Sullivan G, Sherbourne C, Bystritsky A, Stein MB, Roy-Byrne PP** (2009). Computer-assisted delivery of cognitive behavioral therapy for anxiety disorders in primary-care settings. *Depression and Anxiety* **26**, 235–242.
- Cuijpers P** (1997). Bibliotherapy in unipolar depression. *Journal of Behavior Therapy and Experimental Psychiatry* **28**, 139–147.
- Cuijpers P, Schuurmans J** (2007). Self-help interventions for anxiety disorders: an overview. *Current Psychiatry Reports* **9**, 284–290.
- Cuijpers P, van Straten A, Warmerdam L, Andersson G** (2008). Psychological treatment of depression: a meta-analytic database of randomized studies. *BMC Psychiatry* **8**, 36.
- Donner L, Guerney BG** (1969). Automated group desensitization for test anxiety. *Behaviour Research and Therapy* **7**, 1–13.
- Duval S, Tweedie R** (2000). Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* **56**, 455–463.
- Floyd M, Scogin F, McKendree-Smith NL, Floyd DL, Rokke PD** (2004). Cognitive therapy for depression: a comparison of individual psychotherapy and bibliotherapy for depressed older adults. *Behavior Modification* **28**, 297–318.
- Gellatly J, Bower P, Hennessy S, Richards D, Gilbody S, Lovell K** (2007). What makes self-help interventions effective in the management of depressive symptoms? Meta-analysis and meta-regression. *Psychological Medicine* **37**, 1217–1228.
- Ghosh A, Marks IM, Carr A** (1988). Therapist contact and outcome of self-exposure treatment for phobias. *British Journal of Psychiatry* **152**, 234–238.
- Gould RA, Clum GA, Shapiro D** (1993). The use of bibliotherapy in the treatment of panic: a preliminary investigation. *Behavior Therapy* **24**, 241–252.
- Gregory R, Canning S, Lee T, Wise J** (2004). Cognitive bibliotherapy for depression: a meta-analysis. *Professional Psychology, Research and Practice* **35**, 275–280.
- Hecker JE, Losee MC, Fritzier BK, Fink CM** (1996). Self-directed versus therapist-directed cognitive behavioral treatment for panic disorder. *Journal of Anxiety Disorders* **10**, 253–265.
- Hecker JE, Losee MC, Roberson-Nay R, Maki K** (2004). Mastery of Your Anxiety and Panic and brief therapist contact in the treatment of panic disorder. *Journal of Anxiety Disorders* **18**, 111–126.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG** (2003). Measuring inconsistency in meta-analyses. *British Medical Journal* **327**, 557–560.
- Higgins JPT, Green S** (2008). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.0.1* [updated September 2008] (www.cochrane-handbook.org). The Cochrane Collaboration, 2008.
- Hirai M, Clum GA** (2006). A meta-analytic study of self-help interventions for anxiety problems. *Behavior Therapy* **37**, 99–111.
- Hogan RA, Kirchner JH** (1968). Implosive, eclectic verbal and bibliotherapy in the treatment of fears of snakes. *Behaviour Research and Therapy* **6**, 167–171.
- Kahn M, Baker B** (1968). Desensitization with minimal therapist contact. *Journal of Abnormal Psychology* **73**, 198–200.
- Kay-Lambkin FJ, Baker AL, Lewin TJ, Carr VJ** (2009). Computer-based psychological treatment for comorbid depression and problematic alcohol and/or cannabis use: a randomized controlled trial of clinical efficacy. *Addiction* **104**, 378–388.
- Kiropoulos LA, Klein B, Austin DW, Gilson K, Pier C, Mitchell J, Ciechomski L** (2008). Is Internet-based CBT for panic disorder and agoraphobia as effective as face-to-face CBT? *Journal of Anxiety Disorders* **22**, 1273–1284.
- Knaevelsrud C, Maercker A** (2007). Internet-based treatment for PTSD reduces distress and facilitates the development of a strong therapeutic alliance: a randomized controlled clinical trial. *BMC Psychiatry* **7**, 13.
- Lau J, Elliott MA, Jimenez-Silva J, Kupelnick B, Mosteller F, Chalmers TC** (1992). Cumulative meta-analysis of therapeutic trials for myocardial infarction. *New England Journal of Medicine* **327**, 248–254.
- Lidren DM, Watkins PL, Gould RA, Clum GA, Asterino M, Tulloch HL** (1994). A comparison of bibliotherapy and group therapy in the treatment of panic disorder. *Journal of Consulting and Clinical Psychology* **62**, 865–869.
- Lipsey MW** (1990). *Design Sensitivity; Statistical Power for Experimental Research*. Sage: Newbury Park.
- Marks IM, Kenwright M, McDonough M, Whittaker M, Mataix-Cols D** (2004). Saving clinicians' time by delegating routine aspects of therapy to a computer: a randomised controlled trial in phobia/panic disorder. *Psychological Medicine* **34**, 9–18.
- Marrs RW** (1995). A meta-analysis of bibliotherapy studies. *American Journal of Community Psychology* **23**, 843–870.
- Marshall WL, Presse L, Andrews WR** (1976). A self-administered program for public speaking anxiety. *Behaviour Research and Therapy* **14**, 33–39.
- Mataix-Cols D, Cameron R, Gega L, Kenwright M, Marks IM** (2006). Effect of referral source on outcome with cognitive-behavior therapy self-help. *Comprehensive Psychiatry* **47**, 241–245.
- Menchola M, Arkowitz HS, Burke BL** (2007). Efficacy of self-administered treatments for depression and anxiety. *Professional Psychology, Research and Practice* **38**, 421–429.

- Piaggio G, Elbourne DR, Altman DG, Pocock SJ, Evans SJ** (2006). Reporting of noninferiority and equivalence randomized trials: an extension of the CONSORT statement. *Journal of the American Medical Association* **295**, 1152–1160.
- Rosen GM, Glasgow RE, Barrera Jr. M** (1976). A controlled study to assess the clinical efficacy of totally self-administered systematic desensitization. *Journal of Consulting and Clinical Psychology* **44**, 208–217.
- Schmidt MM, Miller WR** (1983). Amount of therapist contact and outcome in a multidimensional depression treatment program. *Acta Psychiatrica Scandinavica* **67**, 319–332.
- Selmi PM, Klein MH, Greist JH, Sorrell S, Erdman HP** (1990). Computer-administered cognitive-behavioral therapy for depression. *American Journal of Psychiatry* **147**, 51–56.
- Shadish WR, Sweeney RB** (1991). Mediators and moderators in meta-analysis: there's a reason we don't let dodo birds tell us which psychotherapies should have prizes. *Journal of Consulting and Clinical Psychology* **59**, 883–893.
- Sharp DM, Power KG, Swanson V** (2000). Reducing therapist contact in cognitive behaviour therapy for panic disorder and agoraphobia in primary care: global measures of outcome in a randomised controlled trial. *British Journal of General Practice* **50**, 963–968.
- Spek V, Cuijpers P, Nyklíček I, Riper H, Keyzer J, Pop V** (2007). Internet-based cognitive behavior therapy for mood and anxiety disorders: a meta-analysis. *Psychological Medicine* **37**, 319–328.
- Spielmanns GI, Pasek LF, McFall JP** (2007). What are the active ingredients in cognitive and behavioral psychotherapy for anxious and depressed children? A meta-analytic review. *Clinical Psychology Review* **27**, 642–654.
- Tillfors M, Carlbring P, Furmark T, Lewenhaupt S, Spak M, Eriksson A, Westling BE, Andersson G** (2008). Treating university students with social phobia and public speaking fears: Internet delivered self-help with or without live group exposure sessions. *Depression and Anxiety* **25**, 708–717.
- Vestre ND, Judge TJ** (1989). Evaluation of self-administered rational emotive therapy programs for interpersonal anxiety. *Journal of Rational-Emotive and Cognitive-Behavior Therapy* **7**, 141–154.
- Watkins PL, Clum GA** (Eds.) (2008). *Handbook of Self-help Therapies*. Routledge: New York.
- Wollersheim JP, Wilson GL** (1991). Group treatment of unipolar depression: a comparison of coping, supportive, bibliotherapy, and delayed treatment groups. *Professional Psychology, Research and Practice* **22**, 496–502.

Appendix A. All effect sizes included from studies comparing guided self-help with face-to-face psychotherapy for depression and anxiety disorders



AQ, Acrophobia Questionnaire; BAI, Beck Anxiety Inventory; BAT, Behavior Approach Test; BCL; Timed Behavioral Checklist for Performance Anxiety; BDI, Beck Depression Inventory; BDI-II, Beck Depression Inventory, 1996 Revision; Blind assess, Blind assessor; BSQ, Body Sensations Questionnaire; CES-D, Center for Epidemiological Studies – Depression scale; CI, confidence interval; clin, clinician; CR, clinician rating of depression; DASS, Depression, Anxiety, Stress Scales; DPAR, Daily Panic Attack Record; FNE, Fear of Negative Evaluation Scale; FQ, Fear Questionnaire; FSS, Fear Survey Schedule;

FT, Fear Thermometer; FTF, face-to-face therapy; G, group; GSH, guided self-help; GSS, global symptom severity; HAMD, Hamilton Rating Scale of Depression; HSCL anx, Hopkins Symptom Checklist, anxiety subscale; I, individual; IG, large group; LSAS, Liebowitz Social Anxiety Scale self-report version; MI, Mobility Inventory; MI-ac, Mobility Inventory (avoidance when accompanied by a trusted person); MI-al, Mobility Inventory (avoidance when alone); MMPI, Minnesota Multiphasic Personality Inventory; MMPI-d, Minnesota Multiphasic Personality Inventory Depression Scale; PA, panic attack; PDSS, Panic Disorder Severity Scale; POMS, Profile of Mood States; SAD, Social Avoidance and Distress Scale; SCL-90-d, Symptom Checklist-90 depression scale; SDS, Self-rating Depression Scale; sG, small group; SIAS, Social Interaction Anxiety Scale; SNAQ, Snake Attitude Questionnaire; SPS, Social Phobia Scale; Std diff, standardized difference; SPSQ, Social Phobia Screening Questionnaire; SR, self-rating of depression; SUD, Subjective Units of Disturbance.